



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$\begin{aligned}
 & -\frac{7}{3}e\frac{\pi}{n}\sin(\zeta + 2\tau + \hat{\xi}) \\
 & + \frac{15}{16}e\frac{\pi}{n}\sin(\zeta - 4\tau + \hat{\xi}).
 \end{aligned}$$

[TO BE CONTINUED.]



NOTE ON A UNIQUE PROPERTY OF AN AXISYMMETRIC DETERMINANT OF THE FOURTH ORDER.

By PROF. THOMAS MUIR, Bishopton, Scotland.

The determinant in question is

$$\begin{vmatrix} 0 & a & b & c \\ a & d & e & f \\ b & e & g & h \\ c & f & h & k \end{vmatrix}, \text{ or } \mathcal{Q} \text{ say,}$$

its only specialty as an axisymmetric determinant of the fourth order being that it has a zero in the place (1,1). Now if we take the minor

$$\begin{vmatrix} 0 & a & b \\ a & d & e \\ b & e & g \end{vmatrix}, \text{ or } K \text{ say,}$$

and substitute in it for a and b the complementary minors of h and f in \mathcal{Q} , it is found that, curiously enough, the new determinant is divisible by the old, the quotient being

$$\begin{vmatrix} 0 & a & b & -c \\ a & d & e & f \\ b & e & g & h \\ c & f & h & 0 \end{vmatrix}$$

That is to say, if H and F denote the complementary minors of h and f in \mathcal{Q} we assert the identity

$$\begin{vmatrix} 0 & H & F \\ H & d & e \\ F & e & g \end{vmatrix} = \begin{vmatrix} 0 & a & b \\ a & d & e \\ b & e & g \end{vmatrix} \cdot \begin{vmatrix} 0 & a & b & -c \\ a & d & e & f \\ b & e & g & h \\ c & f & h & 0 \end{vmatrix}.$$

This may be established in various ways, but perhaps the following is as neat as any other :—

Multiplying the left-hand member column-wise by

$$\begin{vmatrix} 1 & 0 & 0 \\ bc & 1 & 0 \\ -ac & 0 & 1 \end{vmatrix}$$

and bearing in mind that

$$H = \begin{vmatrix} 0 & a & b \\ a & d & e \\ c & f & h \end{vmatrix} = abf + acc - bcd - a^2h,$$

and

$$F = \begin{vmatrix} 0 & a & b \\ b & c & g \\ c & f & h \end{vmatrix} = b^2f + acg - bcc - abh,$$

we have

$$\begin{vmatrix} 0 & H & F \\ H & d & e \\ F & e & g \end{vmatrix} = \begin{vmatrix} bcH - acF & a(bf - ah) & b(bf - ah) \\ H & d & e \\ F & e & g \end{vmatrix}.$$

Multiplying again by the same multiplier, but now performing the operation row-wise in order that the determinant may re-assume the original symmetrical form, we have

$$\begin{vmatrix} 0 & H & F \\ H & d & e \\ F & e & g \end{vmatrix} = \begin{vmatrix} bcH - acF & a(bf - ah) & b(bf - ah) \\ a(bf - ah) & d & e \\ b(bf - ah) & e & g \end{vmatrix}.$$

But since

$$\begin{vmatrix} 0 & a & b & 0 \\ a & d & e & a \\ b & c & g & b \\ c & f & h & c \end{vmatrix} = 0,$$

it follows that

$$cK - bH + aF = 0,$$

and thus

$$bcH - acF = c^2K.$$

Substituting this in our last obtained result, and partitioning the determinant into two, we have

$$\begin{vmatrix} c^2K & 0 & 0 \\ 0 & d & e \\ 0 & e & g \end{vmatrix} + \begin{vmatrix} 0 & a(bf - ah) & b(bf - ah) \\ a(bf - ah) & d & e \\ b(bf - ah) & e & g \end{vmatrix},$$

i. e.,

$$c^2K(dg - e^2) + (bf - ah)^2K,$$

i. e.,

$$K[c^2(dg - e^2) + (bf - ah)^2],$$

i. e.,

$$K \begin{vmatrix} 0 & a & b & -c \\ a & d & e & f \\ b & e & g & h \\ c & f & h & 0 \end{vmatrix},$$

as was to be proved.

In connection with the co-factor of K here it is worth while remarking upon the very considerable cancelling effect of altering the sign of one of the non-axial elements in an axisymmetric determinant. If the determinant be of the n^{th} order the number of terms is of course $n!$, or classifying them as they are got when we expand the determinant according to binary products of elements from the first row and first column, we may say that the number of terms is

$$(n-1)! + (n-2)!(n-1) + 2C_{n-1,2}(n-2)!.$$

Now if one of the non-axial elements be altered in sign, say the element in the place $(1, n)$, the number of terms obtained in like manner is

$$(n-1)! + (n-2)!(n-1) + 2C_{n-2,2}(n-2)!.$$

The number in the latter case is thus diminished by

$$2(n-2)!(C_{n-1,2} - C_{n-2,2})$$

i. e.,

$$2(n-2)!(n-2).$$